

PhytoTechnology Laboratories®

Product Information Sheet

Y8565 YEP Broth **Yeast Extract Peptone Broth**

YP Broth Synonyms:

Properties

Form: Powder

Appearance: Cream to Tan

Application: Molecular Biology, Microbiology, Phytopathology

Solubility: Water

Typical Working

25.0 g/L

Concentration:

Storage Temp: 2 – 6°C

Formula (mg/L)

| Peptone from Meat | 10,000 |
|-------------------|--------|
| Sodium Chloride | 5000 |
| Yeast Extract | 10,000 |

Application Notes

YEP Broth is a standard medium base for the maintenance and propagation of Saccharomyces cerevisiae. This medium can be used as a base for Yeast Peptone Dextrose medium (YPD), or supplemented with another carbon source such as sucrose, glycerol, mannitol, etc.

This medium has been used to culture Agrobacterium tumefaciens (Garfinkel et al., 1981) as well as yeasts such as Saccharomyces (Subashini et al., 2011) and Candida (Jindamorakot et al., 2008).

References

Garfinkel DJ, Simpson RB, Ream LW, White FF, Gordon MP, Nester EW (1981) Genetic analysis of crown gall: fine structure map of the T-DNA by site-directed mutagenesis. Cell 27: 143–153.

Jindamorakot, S., Limtong, S., Yongmanitchai, W., Tuntirungkij, M., Potacharoen, W., Kawasaki, H., Tanticharoen, M. and Nakase, T. (2008), Candida ratchasimensis sp. nov. and Candida khaoyaiensis sp. nov., two anamorphic yeast species isolated from flowers in Thailand. FEMS Yeast Research, 8: 955-960. doi: 10.1111/j.1567-1364.2008.00416.x

Subashini, D., Ejilane, J., Radha, A., Jayasri, M. A., & Suthindhiran, K. (2011). Ethanol production from Sago waste using Saccharomyces cerevisiae Vits-M1. Curr Res J Biol Sci, 3(1), 42-51.

India Contact